

## Iberian Pyrite Belt Subsurface Life (IPBSL), a drilling project in a geochemical Mars terrestrial analogue

**R. Amils** (1, 2), D. Fernández-Remolar (1), V. Parro (1), J.A. Manfredi (1), K. Timmis (3), M. Oggerin (1), M. Sánchez-Román (1), F. J. López (1), J. P. Fernández (1), E. Omoregie (1), D. Gómez-Ortiz (4), C. Briones (1), F. Gómez (1), M. García (1), N. Rodríguez (1), J.L. Sanz (5) and the IPBSL Team.

(1) Centro de Astrobiología (CSIC-INTA), Torrejón de Ardoz, Spain, (2) Centro de Biología Molecular Severo Ochoa (CSIC-UAM), Cantoblanco, Spain, (3) Technical University of Braunschweig, Germany, (4) Universidad Rey Juan Carlos, Madrid, Spain, (5) Departamento de Biología Molecular, Universidad Autónoma de Madrid, Cantoblanco, Spain  
([ramils@cbm.uam.es](mailto:ramils@cbm.uam.es))

### Abstract

Iberian Pyrite Belt Subsurface Life (IPBSL) is a drilling project specifically designed to characterize the subsurface ecosystems operating in the Iberian Pyrite Belt (IPB), in the area of Peña de Hierro, and responsible of the extreme acidic conditions existing in the Rio Tinto basin [1]. Rio Tinto is considered a good geochemical terrestrial analogue of Mars [2, 3]. A dedicated geophysical characterization of the area selected two drilling sites (4) due to the possible existence of water with high ionic content (low resistivity). Two wells have been drilled in the selected area, BH11 and BH10, of depths of 340 and 620 meters respectively, with recovery of cores and generation of samples in anaerobic and sterile conditions. Preliminary results showed an important alteration of mineral structures associated with the presence of water, with production of expected products from the bacterial oxidation of pyrite (sulfates and ferric iron). Ion chromatography of water soluble compounds from uncontaminated samples showed the existence of putative electron donors (ferrous iron, nitrite in addition of the metal sulfides), electron acceptors (sulfate, nitrate, ferric iron) as well as variable concentration of metabolic organic acids (mainly acetate, formate, propionate and oxalate), which are strong signals of the presence of active subsurface ecosystem associated to the high sulfidic mineral content of the IPB. The system is driven by oxidants that appear to be provided by the rock matrix, only groundwater is needed to launch microbial metabolism. The geological, geomicrobiological and molecular biology analysis which are under way, should allow the characterization of this ecosystem of paramount interest in the design of an astrobiological underground Mars exploration mission in the near future.

### Acknowledgements

The work was supported by an ERC grant (contract 250350-IPBSL)

### References

- [1] González-Toril, E., Llobet-Brossa, E., Casamayor, E.O., Amann, R., Amils, R. Microbial ecology of an extreme acidic environment, the Tinto River. *Appl. Environ. Microbiol.*, 69, pp. 4853-4865, 2003
- [2] Fernández-Remolar, D., Morris, R.V., Gruener, J.E., Amils, R., Knoll, A.H. The Rio Tinto Basin, Spain: mineralogy, geobiology and implications for interpretation of outcrop rocks at Meridiani Planum, Mars. *Earth Planet. Sci. Lett.*, 240, pp. 149-167, 2005
- [3] Amils, R., González-Toril, E., Fernández-Remolar, D., Gómez, F., Aguilera, A., Rodríguez, N., Malki, M., García-Moyano, A., González-Fairén, A., de la Fuente, V., Sanz, J.L. Extreme environments as Mars terrestrial analogs: the Rio Tinto case. *Planet. Space Sci.*, 55, pp. 370-381, 2007
- [4] Fernández-Remolar, D. C., Sánchez-Román M., Prieto-Ballesteros, O., Gómez-Ortíz, D., García-Villadangos, M., Molina, A., Rodríguez, N., Barragán, T., Gleenson, D., Banerjee, N., Southam, G., Izawa, M., Loiselle, L., Schmitt-Kopplin, P., Granda, A., Quesada, C., Amils, R. Searching for Biosignatures in Deep Regions of the Rio Tinto Aquifer: The Iberian Pyritic Belt Subsurface Life (IPBSL) Project. *AbSciCon 2012, Atlanta* [#3108]